

KD-Validated Anti-IRAK3 Rabbit Monoclonal Antibody

Rabbit monoclonal antibody


ABG1959

Product Overview


| | |
|-----------------------------------|--|
| Name | KD-Validated Anti-IRAK3 Rabbit Monoclonal Antibody |
| Catalog # | ABG1959 |
| Clonality | Monoclonal |
| Accession(Primary) | Q9Y616 |
| Application Note (Approx.) | WB1:5,000 FC1:2,000 ICC1:1,000 |
| Precautions | |

Target information(Q9Y616)


Synonyms**Gene ID****Other Names****Function****Cellular location****Note**




Western blotting analysis using anti-IRAK3 antibody (Cat#ABG1959). Total cell lysates (30 µg) from various cell lines were loaded and separated by SDS-PAGE. The blot was incubated with anti-IRAK3 antibody (Cat#ABG1959, 1:5,000) and HRP-conjugated goat anti-rabbit secondary antibody respectively.



Western blotting analysis using anti-IRAK3 antibody (Cat#ABG1959). IRAK3 expression in wild type (WT) and IRAK3 shRNA knockdown (KD) HT-1080 cells with 30 μ g of total cell lysates. β -Tubulin serves as a loading control. The blot was incubated with anti-IRAK3 antibody (Cat#ABG1959, 1:5,000) and HRP-conjugated goat anti-rabbit secondary antibody respectively.



Flow cytometric analysis of IRAK3 in HAP-1 cells using IRAK3 antibody (Cat#ABG1959, 1:2,000). Green, isotype control; red, IRAK3.



Immunocytochemical staining of HAP-1 cells with IRAK3 antibody (Cat#ABG1959, 1:1,000). Nuclei were stained blue with DAPI; IRAK3 was stained magenta with Alexa Fluor® 647. Images were taken using Leica stellaris 5. Protein abundance based on laser Intensity and smart gain: Medium. Scale bar: 20 μ m.